

Manganese Catalyzed Desaturation of N-acyl Amines and Ethers

Gang Li, Patrick Kates, Andrew K. Dilger, Peter T.W. Cheng, William R. Ewing, and John T. Groves

ACS Catal., **Just Accepted Manuscript** • DOI: 10.1021/acscatal.9b03457 • Publication Date (Web): 18 Sep 2019

Downloaded from pubs.acs.org on September 23, 2019

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Manganese Catalyzed Desaturation of N-acyl Amines and Ethers

Gang Li,[†] Patrick A. Kates,[†] Andrew K. Dilger,[‡] Peter T. Cheng,[‡] William R. Ewing,[‡] and John T. Groves^{*†}

[†]Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States

[‡]Bristol-Myers Squibb, P. O. Box 5400, Princeton, New Jersey 08543-5400, United States

homogenous catalysis, C-H activation, high-valent metal-oxo complexes, manganese porphyrins, desaturation

ABSTRACT: Enamines and enol ethers are versatile synthons for chemical synthesis. While several methods have been developed to access such molecules, prefunctionalized starting materials are usually required and direct desaturation methods remain rare. Herein, we report direct desaturation reactions of cyclic amines and cyclic ethers using a mild I(III) oxidant, PhI(OAc)₂, and an electron-deficient manganese pentafluorophenylporphyrin catalyst, Mn(TPFPP)Cl. This system displays high efficiency for α,β -desaturation of various cyclic amines and ethers. Mechanistic probes suggest that the desaturation reaction occurs via an initial α -C-H hydroxylation pathway, which serves to protect the product from over-oxidation.


Olefins occupy a key nexus in organic synthesis and process chemistry.¹ Among them, functionalized olefins such as enamines and enol ethers have attracted considerable attention. Enamines are widely represented in natural and synthetic compounds possessing useful biological and physiological properties.²⁻⁷ Enol ethers are used as cross-reaction partners in olefin metathesis,⁸ as well as substrates for access to a variety of functional polymers.⁹⁻¹⁰ In view of this level of utility, numerous methods have been developed to access such olefins. Typically, these methods require prefunctionalized starting materials, thus limiting their practicality.^{2-3, 11-19} Direct, catalytic dehydrogenation would be ideal for the conversion of simple aliphatic starting materials into valuable functionalized olefins, but such methods remain rare. Very recently, Gevorgyan et al. reported a palladium-catalyzed desaturation of aliphatic amines via a hydrogen atom transfer mechanism.²⁰ While efficient for a variety of substrates, the method requires N-benzoyl protecting groups, thus limiting its application due to the forcing conditions required for the removal of that protecting group. Moreover, the method has so far been limited to amines. A more general desaturation method that is applicable to both amines and ethers would be highly valuable.

Our long-standing interest in aliphatic C-H functionalization led us to examine manganese porphyrins and heteroatom rebound catalysis as an approach to amine and ether desaturation.²¹⁻²⁸ In an attempt to functionalize the α -C-H bond of a protected proline (**1a**) using a manganese(III) pentafluorophenyl porphyrin, Mn(TPFPP)Cl, we observed α,β -desaturation with iodosylbenzene as the oxidant (Table 1, entry 1). Although desaturations are commonly observed in aliphatic C-H bond oxidation reactions,²⁹⁻³⁰ this is usually

a minor pathway and olefin oxygenation typically dominates. Thus, this unexpected observation encouraged us to explore the scope of this manganese-catalyzed, direct desaturation protocol.

We initiated this study by screening N-protecting groups using proline derivatives as model substrates. Each substrate resulted solely in desaturation products (Table 1, entries 1-5). Less electron-withdrawing protecting groups (Boc, Bz, Ac) gave better conversions, while more electron-withdrawing protecting groups (Ns, TFA) suppressed the reaction.

Table 1. Manganese-catalyzed desaturation of N-protected proline methyl esters with various oxidants.^a



entry	R	oxidant	conversion	yield
1	Boc	PhIO	35%	30%
2	TFA	PhIO	trace	trace
3	Bz	PhIO	34%	32%
4	Ac	PhIO	28%	25%
5	Ns	PhIO	trace	trace
6	Boc	PhI(OAc) ₂	>95%	>95%
7	Boc	PhI(TFA) ₂	trace	trace
8	Boc	PhI(OPiv) ₂	12%	10%
9	Boc	MesI(OAc) ₂	70%	67%
10	Boc	H ₂ O ₂ /AcOH	trace	trace


11	Boc	<i>m</i> CPBA	trace	trace
12	Cbz	PhI(OAc) ₂	>95%	>95%

^aConditions: substrate (0.1 mmol), PhI(OAc)₂ (0.2 mmol), Mn(TPFPP)Cl (0.005 mmol), MeCN (0.5 mL), 50 °C, 2 h. Conversions and yields determined by GC-MS using naphthalene as an internal standard. Ns = 4-nitrophenylsulfonyl, TFA = trifluoroacetate, Bz = benzoyl, Ac = acetyl, Boc = ^tButyloxycarbonyl, Cbz = carboxybenzyl. Piv = pivaloyl, Mes = mesityl, *m*CPBA = *m*-chloroperoxybenzoic acid.

Carbamate protecting groups such as Boc or Cbz are particularly useful because they can be removed easily.³¹ We then used N-Boc proline methyl ester (**1a**) as the substrate to examine the effect of oxidants in the desaturation reaction. When a milder and more soluble oxidant such as PhI(OAc)₂ was used, we observed quantitative conversion to the desaturated product with no side products (Table 1, entry 6, and Figure S1 in Supporting Information). Switching to a more electrophilic trifluoroacetate oxidant PhI(O₂CCF₃)₂ (Table 1, entry 7) severely suppressed the reaction. The bulkiness of the oxidant was also found to affect the observed reactivity. When PhI(OPiv)₂ was employed as the oxidant (Table 1, entry 8), the desired product was obtained in only 10% yield. A less oxidizing mesityl iodine also resulted in a decreased yield (Table 1, entry 9). Other oxidants that have been previously used in transition-metal-catalyzed proline oxidation³² showed no reactivity under these reaction conditions (Table 1, entries 10 and 11). Cbz-protected proline methyl ester also showed superior reactivity and a quantitative conversion was observed (Table 1, entry 12). A more electron-rich manganese catalyst, Mn(TMP)Cl, showed little catalytic activity (Table S3, entry 2, optimization reactions are presented in the Supporting Information).

We were pleased to find the desaturation reaction proceeded in good yields even with low catalyst loading (Table 2). With 1 mol% catalyst, the reaction was completed in 2 h. Decreasing the catalyst loading to 0.5% still gave a yield of 90%, although a longer reaction time was required. Further decreasing the catalyst loading to 0.1% led to a drastic decrease in yield, possibly due to catalyst-bleaching. No reaction was observed without the catalyst.

Table 2. The effect of catalyst loading on the desaturation reaction.^a



entry	catalyst loading	reaction time	conversion	yield
1	5%	2h	>95%	>95%
2	2%	2h	>95%	>95%
3	1%	2h	>95%	>95%
4	0.5%	4h	95%	90%
5	0.1%	24h	16%	12%
6	no catalyst	48h	<5%	0%

^aConditions: substrate (0.1 mmol), PhI(OAc)₂ (0.2 mmol), MeCN (0.5 mL), 50 °C, 2 h. Conversions and yields determined by GC-MS using naphthalene as an internal standard.

Inspired by the results for proline desaturation, we further investigated the ability of Mn(TPFPP)Cl to catalyze the α,β -desaturation of other amines (Table 3).


Generally, 5- and 6-membered cyclic amines reacted efficiently, affording the desired desaturation products in moderate to excellent yields (**2a-2f**). Morpholine **1g** was also reactive, producing the desired desaturation product **2g** in good yield. It is interesting to note that when bis-N-Boc piperazine **1h** was subjected to the reaction conditions, a double desaturation product **2h** was observed. Similar double desaturation was observed for this substrate with palladium-catalyzed desaturation.²⁰ Notably, substrate reactivity decreased significantly with increasing ring size. Reaction of seven-membered rings (**1i** and **1j**) resulted in moderate yields (**2i** and **2j**), while reaction with a substrate with an eight-membered ring (**1k**) resulted in less than a 20% yield. A non-cyclic amine **1l** was also tested, which only afforded trace desaturation product **2l**. We expect that the transition state for desaturation requires a conformation wherein the incipient C=C bond is in conjugation with the amide group. Such a conformation is readily achieved in small ring substrates but would be more challenging or entropically disfavored in larger ring or acyclic substrates, leading to a lower yield of the enamide product.

Table 3. α,β -Desaturation of amines catalyzed by Mn(TPFPP)Cl.^a


<td> </td>	
<td> </td>	
<td>c</td>	c

^aConditions: substrate (0.2 mmol), PhI(OAc)₂ (0.4 mmol), Mn(TPFPP)Cl (0.002 mmol), MeCN (1 mL), 50 °C, 2 h, isolated yield reported unless otherwise stated. ^b0.5 mmol scale. ^cConversion and yield determined by GC-MS.

Considering the electronic similarities between cyclic amines and cyclic ethers, we applied this desaturation reaction to a panel of cyclic ether structures. Indeed, when tetrahydropyran **1m** was subjected to the desaturation conditions with a higher catalyst loading, the desired product **2m** was observed in 55% NMR yield (eq. 3).




Similarly, the tetrahydropyran ester **1n** afforded only the desaturated product **2n**, which was isolated in 62% yield (eq. 4).



Owing to the fact that α,β -unsaturated cyclic en-carbamates are versatile intermediates for the synthesis of various nitrogen-containing bioactive molecules,^{6, 33-37} it would be highly advantageous if the viability of the desaturation reaction was demonstrated on a larger scale. Accordingly, we then carried out a gram-scale reaction on **1a** to explore reaction scalability. The reaction worked smoothly with no significant decrease in yield (Table 3, condition b for **2a**).

We carried out the reaction under air-free conditions in dry solvent to gain some mechanistic insight. No conversion was observed under these conditions even after heating for 24 h. However, upon addition of 10 equiv of water, the desaturated product quickly appeared in quantitative yield (eq. 5). We attribute this water accelerating effect to slow hydrolysis of $\text{PhI}(\text{OAc})_2$ in wet solvent to form PhIO or $\text{PhI}(\text{OH})(\text{OAc})$, which has also been observed in other metalloporphyrin-catalyzed oxidation reactions using $\text{PhI}(\text{OAc})_2$ as oxidant.³⁸⁻³⁹



Examination of the reaction mixture by UV-Vis spectroscopy was also informative. The $\text{Mn}(\text{TPFPP})\text{Cl}$ catalyst displayed a split Soret band at 359 and 470 nm and a Q band at 566 nm in acetonitrile, typical for manganese(III) porphyrin complexes. After addition of $\text{PhI}(\text{OAc})_2$ (10 equiv), the band at 470 nm disappeared and a new band at 418 nm was formed. At the same time, the Q band at 566 nm also faded while a new band was observed at 540 nm (Figure 1). The new peaks match previous reported values for oxo- Mn^{IV} species.⁴⁰⁻⁴³ Zhang et al. have reported similar spectra in a previous study on oxidation of alkenes and activated benzylic hydrocarbons.³⁹







Figure 1. UV-Vis spectra observed upon addition of $\text{PhI}(\text{OAc})_2$ to $\text{Mn}(\text{TPFPP})\text{Cl}$: a) Soret region; b) Q band region. Solid line: $\text{Mn}(\text{TPFPP})\text{Cl}$ (10 μM) in acetonitrile; dotted line: 30 s after the addition of $\text{PhI}(\text{OAc})_2$ (20 equiv).

The reactivity of the $\text{Mn}(\text{TPFPP})\text{Cl}/\text{PhI}(\text{OAc})_2$ system indicates the involvement of an oxo- Mn^{IV} intermediate.^{28, 39} The observed oxo- Mn^{IV} species is likely to be generated upon rapid comproportionation of oxo- Mn^{V} with a Mn^{III} species⁴³ and is unreactive toward these N-protected amines.

Oxo- Mn^{V} porphyrins have been shown to oxidize olefins rapidly.^{26, 44-47} Also, $\text{PhI}(\text{OAc})_2$ is known to react with enamines.⁴⁸ Thus, we were curious as to why the desaturated products could be isolated in such high yields. Interestingly, when a reaction mixture with the N-benzoyl substrate **1c** was examined by mass spectrometry, a hydroxylated product (presumably the hemi-aminal) was observed (see Supporting Information). This result is analogous to observations by White with a non-heme iron catalyst.³² Bietti and Costas have also reported similar α -hydroxylations of amines using a non-heme manganese catalyst.⁴⁹⁻⁵² This unstable hemi-aminal product, or its acyloxy derivative, is presumably converted rapidly into the desired desaturated product during work-up. No hydroxylated intermediate was observed for the corresponding N-Boc protected substrate **1a**, possibly due to the greater instability of the corresponding hemi-aminals. This hemi-aminal intermediate is likely converted to an acyloxy-aminal intermediate in the presence of excess $\text{PhI}(\text{OAc})_2$, which could suppress further oxidation by the reactive oxo- Mn^{V} species due to its steric hindrance and electron-deficiency. In contrast, when the desaturated product **2a** was directly subjected to this $\text{Mn}(\text{TPFPP})\text{Cl}/\text{PhI}(\text{OAc})_2$ system, it was further oxidized and no substrate was recovered. Interestingly, when the cyclic lactam substrate **1o** was subjected to this desaturation conditions, the α -hydroxylated product **2o** was isolated in high yield (eq. 6), which also supports the existence of a hemi-aminal intermediate in the desaturation reaction.



In the case of cyclic ether **1m**, the corresponding hemi-acetal is stable. When this hemi-acetal was directly reacted with $\text{PhI}(\text{OAc})_2$ in acetonitrile (eq 7), the desaturated product was observed, indicating that hemi-acetal could indeed be the intermediate of the desaturation reaction.




5
6
7
8
9
10
11
12

In accord with these results, we propose a mechanism for this manganese-catalyzed desaturation reaction (Figure 2). In the presence of oxidant, the manganese(III) porphyrin will be oxidized to an oxo-Mn^V species that abstracts the α -hydrogen atom from the substrate, resulting in a hydro-Mn^{IV} species and a substrate radical. Oxygen rebound of the substrate radical forms the unstable hemiaminal species, which dehydrates to yield the desired desaturated product.

13
14
15
16
17
18
19
20
21

Desaturation reactions of this type are commonly observed with various heme⁵³ and non-heme iron⁵⁴ oxidases. It is noteworthy that assistance from the adjacent heteroatom in C-C desaturation has also been observed in a non-heme Fe/2OG system.⁵⁴ However, a hydroxylated intermediate is not usually proposed in the mechanism.⁵⁴ Our result suggests that a hydroxylation pathway may be taken into consideration in desaturations in biological systems.



34
35

Figure 2. Proposed mechanism for manganese-catalyzed desaturation (X = N-PG or O).

36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

In conclusion, we have demonstrated that an electron-deficient manganese porphyrin, Mn(TPFPP)Cl, is a highly efficient catalyst for the α,β -desaturation of various cyclic amines and cyclic ethers to afford a range of highly useful functionalized olefins. The reaction protocol is simple, and the reaction conditions are very mild and operationally simple. The homogeneous conditions and tolerance of air and moisture make this protocol amenable to automated and high through-put procedures. We have also shown that this reaction has the potential to be applied at larger-scale. Our preliminary mechanistic studies indicate that the reaction proceeds via a C-H hydroxylation-dehydration pathway. An intermediate hemiaminal for cyclic amine substrate is resistant to further oxidation, resulting in a highly efficient desaturation. Further extensions of this reaction are currently ongoing.

51 52 ASSOCIATED CONTENT

53 Supporting Information

54
55

The Supporting Information is available free of charge on the ACS Publications website.

Detailed condition optimization; experimental procedures for desaturation; and characterization data, including ¹H and ¹³C NMR spectra (PDF)

56 AUTHOR INFORMATION

57 Corresponding Author

*E-mail: jtgroves@princeton.edu

58 ORCID

John T. Groves: 0000-0002-9944-5899

Gang Li: 0000-0001-6680-961X

59 Notes

The authors declare no competing financial interest.

60 ACKNOWLEDGMENT

This research was supported by the US National Science Foundation awards CHE-1464578 and CHE-1900048 (JTG) and the BMS Center for Molecular Synthesis at Princeton University.

REFERENCES

- (1) Zimmermann, H.; Walzl, R., *Ullman's Encyclopedia of Industrial Chemistry*. Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2009.
- (2) Cook, A. G., *Enamines: Synthesis, Structure, and Reactions*. 2 ed.; Marcel Dekker, Inc.: New York, 1988.
- (3) Rappoport, Z., *The Chemistry of Enamines*. Wiley & Sons: New York, 1994.
- (4) Pan, W. D.; Lahue, B. R.; Ma, Y.; Nair, L. G.; Shipps, G. W.; Wang, Y. L.; Doll, R.; Bogen, S. L., Core Modification of Substituted Piperidines as Novel Inhibitors of HDM2-p53 Protein-Protein Interaction. *Bioorg. Med. Chem. Lett.* **2014**, *24*, 1983-1986.
- (5) Elliott, M. L.; Thomas, K.; Kennedy, S.; Koduri, N. D.; Hussaini, R. S.; Sheaff, R. J., Identification of Novel Proteasome Inhibitors from an Enaminone Library. *Chem. Biol. Drug Des.* **2015**, *86*, 322-332.
- (6) Steele, A. D.; Keohane, C. E.; Knouse, K. W.; Rossiter, S. E.; Williams, S. J.; Wuest, W. M., Diverted Total Synthesis of Promysalin Analogs Demonstrates That an Iron-Binding Motif Is Responsible for Its Narrow-Spectrum Antibacterial Activity. *J. Am. Chem. Soc.* **2016**, *138*, 5833-5836.
- (7) Kaduskar, R. D.; Dhavan, A. A.; Dallavalle, S.; Scaglioni, L.; Musso, L., Total Synthesis of the Saticyldehydroproline-Containing Antibiotic Promysalin. *Tetrahedron* **2016**, *72*, 2034-2041.
- (8) Hoveyda, A. H., Evolution of Catalytic Stereoselective Olefin Metathesis: From Ancillary Transformation to Purveyor of Stereochemical Identity. *J. Org. Chem.* **2014**, *79*, 4763-4792.
- (9) Sugihara, S.; Hashimoto, K.; Matsumoto, Y.; Kanaoka, S.; Aoshima, S., Thermosensitive polyalcohols: Synthesis via Living Cationic Polymerization of Vinyl Ethers with a Silyloxy Group. *J. Polym. Sci. Pol. Chem.* **2003**, *41*, 3300-3312.
- (10) Tracton, A. A., *Coating Materials and Surface Coatings*. CRC Press: Boca Raton, FL, 2006.
- (11) Hickmott, P. W., Enamines - Recent Advances in Synthetic, Spectroscopic, Mechanistic, and Stereochemical Aspects, *Tetrahedron* **1982**, *38*, 1975-2050.
- (12) Hickmott, P. W., Enamines - Recent Advances in Synthetic, Spectroscopic, Mechanistic, and Stereochemical Aspects, *Tetrahedron* **1982**, *38*, 3363-3446.
- (13) Oliveira, D. F.; Miranda, P. C. M. L.; Correia, C. R. D., Efficient and Expedient Protocols for the Synthesis of Racemic and Enantiomerically Pure Endocyclic Enecarbamates from N-Acyl Lactams and N-Acyl Pyrrolidines. *J. Org. Chem.* **1999**, *64*, 6646-6652.
- (14) Pohlki, F.; Doye, S., The Catalytic Hydroamination of Alkynes. *Chem. Soc. Rev.* **2003**, *32*, 104-114.
- (15) Yu, J. R.; Truc, V.; Riebel, P.; Hierl, E.; Mudryk, B., One-pot Synthesis of Cyclic Enecarbamates from Lactam Carbamates. *Tetrahedron Lett.* **2005**, *46* (23), 4011-4013.

- (16) Dehli, J. R.; Legros, J.; Bolm, C., Synthesis of Enamines, Enol Ethers and Related Compounds by Cross-Coupling Reactions. *Chem. Commun.* **2005**, 973-986.
- (17) Severin, R.; Doye, S., The Catalytic Hydroamination of Alkynes. *Chem. Soc. Rev.* **2007**, *36*, 1407-1420.
- (18) Yu, Y. F.; Shu, C.; Zhou, B.; Li, J. Q.; Zhou, J. M.; Ye, L. W., Efficient and Practical Synthesis of Enantioenriched 2,3-Dihydropyrroles through Gold-Catalyzed anti-Markovnikov Hydroamination of Chiral Homopropargyl Sulfonamides. *Chem. Commun.* **2015**, *51*, 2126-2129.
- (19) Patel, M.; Saunthwal, R. K.; Verma, A. K., Base-Mediated Hydroamination of Alkynes. *Acc. Chem. Res.* **2017**, *50* (2), 240-254.
- (20) Chuentragool, P.; Parasram, M.; Shi, Y.; Gevorgyan, V., General, Mild, and Selective Method for Desaturation of Aliphatic Amines. *J. Am. Chem. Soc.* **2018**, *140*, 2465-2468.
- (21) Liu, W.; Huang, X.; Cheng, M. J.; Nielsen, R. J.; Goddard, W. A.; Groves, J. T., Oxidative Aliphatic C-H Fluorination with Fluoride Ion Catalyzed by a Manganese Porphyrin. *Science* **2012**, *337*, 1322-1325.
- (22) Huang, X.; Bergsten, T. M.; Groves, J. T., Manganese-Catalyzed Late-Stage Aliphatic C-H Azidation. *J. Am. Chem. Soc.* **2015**, *137*, 5300-5303.
- (23) Liu, W.; Groves, J. T., Manganese Catalyzed C-H Halogenation. *Acc. Chem. Res.* **2015**, *48*, 1727-1735.
- (24) Liu, W.; Cheng, M. J.; Nielsen, R. J.; Goddard, W. A.; Groves, J. T., Probing the C-O Bond-Formation Step in Metalloporphyrin-Catalyzed C-H Oxygenation Reactions. *ACS Catal.* **2017**, *7*, 4182-4188.
- (25) Huang, X.; Zhuang, T.; Kates, P. A.; Gao, H.; Chen, X.; Groves, J. T., Alkyl Isocyanates via Manganese-Catalyzed C-H Activation for the Preparation of Substituted Ureas. *J. Am. Chem. Soc.* **2017**, *139*, 15407-15413.
- (26) Huang, X.; Groves, J. T., Oxygen Activation and Radical Transformations in Heme Proteins and Metalloporphyrins. *Chem. Rev.* **2018**, *118*, 2491-2553.
- (27) Liu, W.; Huang, X.; Placzek, M. S.; Krska, S. W.; McQuade, P.; Hooker, J. M.; Groves, J. T., Site-Selective ¹⁸F-Fluorination of Unactivated C-H Bonds Mediated by a Manganese Porphyrin. *Chem. Sci.* **2018**, *9*, 1168-1172.
- (28) Li, G.; Dilger, A. K.; Cheng, P. T.; Ewing, W. R.; Groves, J. T., Selective C-H Halogenation with a Highly Fluorinated Manganese Porphyrin. *Angew. Chem. Int. Ed.* **2018**, *57*, 1251-1255.
- (29) Broun, P.; Shanklin, J.; Whittle, E.; Somerville, C., Catalytic Plasticity of Fatty Acid Modification Enzymes Underlying Chemical Diversity of Plant Lipids. *Science* **1998**, *282*, 1315-1317.
- (30) Broadwater, J. A.; Whittle, E.; Shanklin, J., Desaturation and Hydroxylation - Residues 148 and 324 of Arabidopsis FAD2, in Addition to Substrate Chain Length, Exert a Major Influence in Partitioning of Catalytic Specificity. *J. Biol. Chem.* **2002**, *277*, 15613-15620.
- (31) Gibson, F. S.; Bergmeier, S. C.; Rapoport, H., Selective Removal of an N-BOC Protecting Group in the Presence of a tert-Butyl Ester and Other Acid-Sensitive Groups. *J. Org. Chem.* **1994**, *59*, 3216-3218.
- (32) Osberger, T. J.; Rogness, D. C.; Kohrt, J. T.; Stepan, A. F.; White, M. C., Oxidative Diversification of Amino Acids and Peptides by Small-Molecule Iron Catalysis. *Nature* **2016**, *537*, 214-219.
- (33) Matos, M. N.; Afonso, C. A. M.; Batey, R. A., Synthesis of Substituted Pyrrolidines and Piperidines from Endocyclic Enamine Derivatives. Synthesis of (+/-)-Laburnamine. *Tetrahedron* **2005**, *61*, 1221-1244.
- (34) Kaname, M.; Yoshifuji, S.; Sashida, H., An Efficient Transformation of Cyclic Ene-Carbamates into omega-(N-Formylamino)carboxylic Acids by Ruthenium Tetroxide Oxidation. *Chem. Pharm. Bull.* **2008**, *56*, 1310-1313.
- (35) Savage, S. A.; Jones, G. S.; Kolotuchin, S.; Ramrattan, S. A.; Vu, T.; Waltermire, R. E., Preparation of Saxagliptin, a Novel DPP-IV Inhibitor. *Org. Proc. Res. Dev.* **2009**, *13*, 1169-1176.
- (36) Cheng, B.; Wu, F. F.; Yang, X. B.; Zhou, Y. D.; Wan, X. L.; Zhai, H. B., Efficient Total Synthesis of Marine Alkaloid (-)-Nakodamarin A. *Chem. Eur. J.* **2011**, *17*, 12569-12572.
- (37) Ding, D.; Pan, X. H.; Yu, W. S.; Li, X. J.; Chen, S. K.; Liu, F., An Cost-Effective and Safe Process of L-cis-4,5-Methanoproline Amide, the Key Synthetic Intermediate of Saxagliptin, via an Improved Simmons-Smith Reaction. *Heterocycles* **2015**, *91*, 719-725.
- (38) In, J. H.; Park, S. E.; Song, R.; Nam, W., Iodobenzene Diacetate as an Efficient Terminal Oxidant in Iron(III) Porphyrin Complex-Catalyzed Oxygenation Reactions. *Inorg. Chim. Acta* **2003**, *343*, 373-376.
- (39) Kwong, K. W.; Chen, T. H.; Luo, W. L.; Jeddi, H.; Zhang, R., A Biomimetic Oxidation Catalyzed by Manganese(III) Porphyrins and Iodobenzene Diacetate: Synthetic and Mechanistic Investigations. *Inorg. Chim. Acta* **2015**, *430*, 176-183.
- (40) Schappacher, M.; Weiss, R., Formation of Manganese(IV)-Oxo-Porphyrin Derivatives by Decomposition of Peroxycarbonate Complexes. *Inorg. Chem.* **1987**, *26*, 1189-1190.
- (41) Groves, J. T.; Stern, M. K., Synthesis, Characterization, and Reactivity of Oxomanganese(IV) Porphyrin Complexes. *J. Am. Chem. Soc.* **1988**, *110*, 8628-8638.
- (42) Czernuszewicz, R. S.; Su, Y. O.; Stern, M. K.; Macor, K. A.; Kim, D.; Groves, J. T.; Spiro, T. G., Oxomanganese(IV) Porphyrins Identified by Resonance Raman and Infrared-Spectroscopy - Weak Bonds and the Stability of the Half-Filled T_{2g} Subshell. *J. Am. Chem. Soc.* **1988**, *110*, 4158-4165.
- (43) Zhang, R.; Horner, J. H.; Newcomb, M., Laser Flash Photolysis Generation and Kinetic Studies of Porphyrin-Manganese-Oxo Intermediates. Rate Constants for Oxidations Effected by Porphyrin-Mn^V-Oxo Species and Apparent Disproportionation Equilibrium Constants for Porphyrin-Mn^{IV}-Oxo Species. *J. Am. Chem. Soc.* **2005**, *127*, 6573-6582.
- (44) Baglia, R. A.; Zaragoza, J. P. T.; Goldberg, D. P., Biomimetic Reactivity of Oxygen-Derived Manganese and Iron Porphyrinoid Complexes. *Chem. Rev.* **2017**, *117*, 13320-13352.
- (45) Groves, J. T.; Watanabe, Y.; Mcmurry, T. J., Oxygen Activation by Metalloporphyrins - Formation and Decomposition of an Acylperoxymanganese(III) Complex. *J. Am. Chem. Soc.* **1983**, *105*, 4489-4490.
- (46) Groves, J. T.; Watanabe, Y., Heterolytic and Homolytic O-O Bond-Cleavage Reactions of (Acylperoxo)Manganese(III) Porphyrins. *Inorg. Chem.* **1986**, *25*, 4808-4810.
- (47) Jin, N.; Ibrahim, M.; Spiro, T. G.; Groves, J. T., Trans-dioxo Manganese(V) Porphyrins. *J. Am. Chem. Soc.* **2007**, *129*, 12416-12417.
- (48) Chen, M.; Zhang, W.; Ren, Z. H.; Gao, W. Y.; Wang, Y. Y.; Guan, Z. H., PhI(OAc)₂-Promoted Umpolung Acetoxylation of Enamides for the Synthesis of α -Acetoxy Ketones. *Sci. China Chem.* **2017**, *60*, 761-768.
- (49) Milan, M.; Carboni, G.; Salamone, M.; Costas, M.; Bietti, M., Tuning Selectivity in Aliphatic C-H Bond Oxidation of N-Alkylamides and Phthalimides Catalyzed by Manganese Complexes. *ACS Catal.* **2017**, *7*, 5903-5911.
- (50) Milan, M.; Salamone, M.; Costas, M.; Bietti, M., The Quest for Selectivity in Hydrogen Atom Transfer Based Aliphatic C-H Bond Oxygenation. *Acc. Chem. Res.* **2018**, *51*, 1984-1995.
- (51) Dantignana, V.; Milan, M.; Cussó, O.; Company, A.; Bietti, M.; Costas, M., Chemoselective Aliphatic C-H Bond Oxidation Enabled by Polarity Reversal. *ACS Cent. Sci.* **2017**, *3*, 1350-1358.
- (52) Bietti, M., Activation and Deactivation Strategies Promoted by Medium Effects for Selective Aliphatic C-H Bond Functionalization. *Angew. Chem. Int. Ed.* **2018**, *57*, 16618-16637.
- (53) Denisov, I. G.; Makris, T. M.; Sligar, S. G.; Schlichting, I., Structure and Chemistry of Cytochrome P450. *Chem. Rev.* **2005**, *105*, 2253-2277.
- (54) Dunham, N. P.; Chang, W. C.; Mitchell, A. J.; Martinie, R. J.; Zhang, B.; Bergman, J. A.; Rajakovich, L. J.; Wang, B.; Silakov, A.; Krebs, C.; Boal, A. K.; Bollinger, J. M., Two Distinct Mechanisms for C-C Desaturation by Iron(II)- and 2-(Oxo)glutarate-Dependent Oxygenases: Importance of α -Heteroatom Assistance. *J. Am. Chem. Soc.* **2018**, *140*, 7116-7126.

