



## CCAPS National Comparison — Pre-Post Change

VER. 5/2017

Based on 632 unique local clients, with at least 2 CCAPS Administrations from 07/01/2014 to 06/30/2015.  
The clients in the resulting sample have an average of 5 administrations.



This report compares your center's average change on the CCAPS' subscales to a national sample of 106 counseling centers representing 47,948 clients. Average change is calculated by subtracting each client's last scorable CCAPS administration from their first scorable CCAPS administration and then averaging these differences by subscale. Two charts are provided above based on client's level of initial distress: (1) clients with moderate and/or elevated initial distress or (2) clients with only elevated initial distress.

**For example**, looking at the first chart (moderate + elevated), your center's average change for Depression is at the 62nd percentile. This means that your center's average change on the Depression subscale is greater than the change achieved by 62 % of counseling centers in the national sample (for clients who's initial distress was at least moderate and/or elevated).

## CCAPS Pre-Post Change Data Table

Subscale by Initial Distress Level	Local Average Change in Raw Subscale Score	# of Local Clients Above Cut Score	% of Local Clients Above Cut Score	National Average Change in Raw Subscale Score	# of National Clients Above Cut Score	% of National Clients above Cut Score
Depression Low-Cut	0.732	541	85.6 %	0.673	34,356	71.7 %
Depression High-Cut	0.908	403	63.8 %	0.835	24,369	50.8 %
Generalized Anxiety Low-Cut	0.596	479	75.8 %	0.523	35,923	74.9 %
Generalized Anxiety High-Cut	0.765	311	49.2 %	0.684	22,916	47.8 %
Social Anxiety Low-Cut	0.466	454	71.8 %	0.419	30,170	62.9 %
Social Anxiety High-Cut	0.619	283	44.8 %	0.525	17,372	36.2 %
Academic Distress Low-Cut	0.373	459	72.6 %	0.475	32,448	67.7 %
Academic Distress High-Cut	0.673	246	38.9 %	0.776	14,468	30.2 %
Eating Concerns Low-Cut	0.537	231	36.6 %	0.595	16,463	34.3 %
Eating Concerns High-Cut	0.636	173	27.4 %	0.662	13,343	27.8 %
Hostility Low-Cut	0.658	333	52.7 %	0.545	22,581	47.1 %
Hostility High-Cut	0.818	217	34.3 %	0.780	11,604	24.2 %
Alcohol Use Low-Cut	0.618	117	18.5 %	0.570	15,870	33.1 %
Alcohol Use High-Cut	0.741	79	12.5 %	0.749	10,163	21.2 %
Distress Index Low-Cut	0.556	523	82.8 %	0.490	36,138	75.4 %
Distress Index High-Cut	0.693	278	44.0 %	0.672	16,856	35.2 %

### Data Table Column Descriptions

Subscale by Initial Distress Level	Scores on the CCAPS are divided into three ranges of distress, Low-Distress (white), Moderate Distress (yellow), and Elevated Distress (red), using Low and High cut-scores (Low-Cut and High-Cut). Clients that score over the Low-Cut include Moderate Distress plus Elevated Distress scores. Those that score over the High-Cut include Elevated Distress scores only. For more information about cut-scores please view the CCAPS User Manual under "CCAPS" in the Help Tab.
Local Average Change in Raw Subscale Score	The average difference in Pre-Post CCAPS scores at your center for a given subscale and initial distress.
# of Local Clients Above Cut Score	The number of clients at your center that scored above the indicated cut-score (low or high) per subscale.
% of Local Clients Above Cut Score	The percentage of clients at your center that scored the indicated cut-score (low or high) per subscale.
National Average Change in Raw Subscale Score	The national average Pre-Post change in CCAPS scores for the indicated subscale and cut-score.
# of National Clients Above Cut Score	The number of clients in the national sample that scored above the indicated subscale and cut-score.
% of National Clients Above Cut Score	The percentage of clients in the national sample that that scored above the indicated subscale and cut-score.