



## CCAPS National Comparison — Pre-Post Change

VER. 5/2017

Based on 710 unique local clients, with at least 2 CCAPS Administrations from 07/01/2015 to 06/30/2016.  
The clients in the resulting sample have an average of 6 administrations.



This report compares your center's average change on the CCAPS' subscales to a national sample of 106 counseling centers representing 47,948 clients. Average change is calculated by subtracting each client's last scorable CCAPS administration from their first scorable CCAPS administration and then averaging these differences by subscale. Two charts are provided above based on client's level of initial distress: (1) clients with moderate and/or elevated initial distress or (2) clients with only elevated initial distress.

**For example**, looking at the first chart (moderate + elevated), your center's average change for Depression is at the 72nd percentile. This means that your center's average change on the Depression subscale is greater than the change achieved by 72 % of counseling centers in the national sample (for clients who's initial distress was at least moderate and/or elevated).

## CCAPS Pre-Post Change Data Table

Subscale by Initial Distress Level	Local Average Change in Raw Subscale Score	# of Local Clients Above Cut Score	% of Local Clients Above Cut Score	National Average Change in Raw Subscale Score	# of National Clients Above Cut Score	% of National Clients above Cut Score
Depression Low-Cut	0.786	604	85.1 %	0.673	34,356	71.7 %
Depression High-Cut	0.962	447	63.0 %	0.835	24,369	50.8 %
Generalized Anxiety Low-Cut	0.572	546	76.9 %	0.523	35,923	74.9 %
Generalized Anxiety High-Cut	0.713	367	51.7 %	0.684	22,916	47.8 %
Social Anxiety Low-Cut	0.446	517	72.8 %	0.419	30,170	62.9 %
Social Anxiety High-Cut	0.581	316	44.5 %	0.525	17,372	36.2 %
Academic Distress Low-Cut	0.510	522	73.5 %	0.475	32,448	67.7 %
Academic Distress High-Cut	0.769	282	39.7 %	0.776	14,468	30.2 %
Eating Concerns Low-Cut	0.662	295	41.5 %	0.595	16,463	34.3 %
Eating Concerns High-Cut	0.728	232	32.7 %	0.662	13,343	27.8 %
Hostility Low-Cut	0.571	363	51.1 %	0.545	22,581	47.1 %
Hostility High-Cut	0.755	209	29.4 %	0.780	11,604	24.2 %
Alcohol Use Low-Cut	0.859	135	19.0 %	0.570	15,870	33.1 %
Alcohol Use High-Cut	1.030	96	13.5 %	0.749	10,163	21.2 %
Distress Index Low-Cut	0.582	575	81.0 %	0.490	36,138	75.4 %
Distress Index High-Cut	0.727	333	46.9 %	0.672	16,856	35.2 %

### Data Table Column Descriptions

Subscale by Initial Distress Level	Scores on the CCAPS are divided into three ranges of distress, Low-Distress (white), Moderate Distress (yellow), and Elevated Distress (red), using Low and High cut-scores (Low-Cut and High-Cut). Clients that score over the Low-Cut include Moderate Distress plus Elevated Distress scores. Those that score over the High-Cut include Elevated Distress scores only. For more information about cut-scores please view the CCAPS User Manual under "CCAPS" in the Help Tab.
Local Average Change in Raw Subscale Score	The average difference in Pre-Post CCAPS scores at your center for a given subscale and initial distress.
# of Local Clients Above Cut Score	The number of clients at your center that scored above the indicated cut-score (low or high) per subscale.
% of Local Clients Above Cut Score	The percentage of clients at your center that scored the indicated cut-score (low or high) per subscale.
National Average Change in Raw Subscale Score	The national average Pre-Post change in CCAPS scores for the indicated subscale and cut-score.
# of National Clients Above Cut Score	The number of clients in the national sample that scored above the indicated subscale and cut-score.
% of National Clients Above Cut Score	The percentage of clients in the national sample that that scored above the indicated subscale and cut-score.